

A case for weakening the Church-Turing Thesis

Bhupinder Singh Anand

Draft of January 3, 2012.*

Abstract

We conclude from Gödel’s Theorem VII of his seminal 1931 paper that every recursive function $f(x_1, x_2)$ is representable in the first-order Peano Arithmetic PA by a formula $[F(x_1, x_2, x_3)]$ which is algorithmically verifiable, but not algorithmically computable, *if* we assume that the negation of a universally quantified formula of the first-order predicate calculus is always indicative of the existence of a counter-example under the standard interpretation of PA. We conclude that the standard postulation of the Church-Turing Thesis does not hold if we define a number-theoretic formula as effectively computable if, and only if, it is algorithmically verifiable; and needs to be replaced by a weaker postulation of the Thesis as an equivalence.

Keywords Algorithmic computability, algorithmic verifiability, Aristotle’s particularisation, Church-Turing Thesis, effective computability, first-order, Gödel β -function, Peano Arithmetic PA, standard interpretation, Tarski, uniform method.

1 Introduction

We begin by noting that the following theses are classically equivalent¹:

Standard Church’s Thesis² A number-theoretic function (or relation, treated as a Boolean function) is effectively computable if, and only if, it is partial-recursive³.

Standard Turing’s Thesis⁴ A number-theoretic function (or relation, treated as a Boolean function) is effectively computable if, and only if, it is Turing-computable⁵.

In this paper we shall argue that the principle of Occam’s razor suggests the Theses should be postulated minimally as the following equivalences:

*Subject class: LO; MSC: 03B10

¹cf. [Me64], p.237.

²*Church’s (original) Thesis* The effectively computable number-theoretic functions are the algorithmically computable number-theoretic functions [Ch36].

³cf. [Me64], p.227.

⁴After describing what he meant by “computable” numbers in the opening sentence of his 1936 paper on Computable Numbers [Tu36], Turing immediately expressed this thesis—albeit informally—as: “...the computable numbers include all numbers which could naturally be regarded as computable”.

⁵cf. [BBJ03], p.33.

Weak Church’s Thesis A number-theoretic function (or relation, treated as a Boolean function) is effectively computable if, and only if, it is instantiationally equivalent to a partial-recursive function (or relation, treated as a Boolean function).

Weak Turing’s Thesis A number-theoretic function (or relation, treated as a Boolean function) is effectively computable if, and only if, it is instantiationally equivalent to a Turing-computable function (or relation, treated as a Boolean function).

1.1 The need for explicitly distinguishing between ‘instantiation’ and ‘uniform’ methods

It is significant that both Kurt Gödel (initially) and Alonzo Church (subsequently—possibly under the influence of Gödel’s disquietitude) enunciated Church’s formulation of ‘effective computability’ as a Thesis because Gödel was instinctively uncomfortable with accepting it as a definition that *minimally* captures the essence of ‘*intuitive* effective computability’⁶.

Gödel’s reservations seem vindicated if we accept that a number-theoretic function can be effectively computable instantiationally (in the sense of being algorithmically *verifiable* as defined in Section 5 below), but not by a uniform method (in the sense of being algorithmically *computable* as defined in Section 5).

The significance of the fact (considered below in Section 6) that ‘truth’ too can be effectively decidable *both* instantiationally *and* by a uniform (algorithmic) method under the standard interpretation of PA is reflected in Gödel’s famous 1951 Gibbs lecture⁷, where he remarks:

“I wish to point out that one may conjecture the truth of a universal proposition (for example, that I shall be able to verify a certain property for any integer given to me) and at the same time conjecture that no general proof for this fact exists. It is easy to imagine situations in which both these conjectures would be very well founded. For the first half of it, this would, for example, be the case if the proposition in question were some equation $F(n) = G(n)$ of two number-theoretical functions which could be verified up to very great numbers n .”⁸

Such a possibility is also implicit in Turing’s remarks⁹:

“The computable numbers do not include all (in the ordinary sense) definable numbers. Let P be a sequence whose n -th figure is 1 or 0 according as n is or is not satisfactory. It is an immediate consequence of the theorem of §8 that P is not computable. It is (so far as we know at present) possible that any assigned number of figures of P can be calculated, but not by a uniform process. When sufficiently

⁶See [Si97].

⁷[Go51].

⁸Parikh’s paper [Pa71] can also be viewed as an attempt to investigate the consequences of expressing the essence of Gödel’s remarks formally.

⁹[Tu36], §9(II), p.139.

many figures of P have been calculated, an essentially new method is necessary in order to obtain more figures.”

The need for placing such a distinction on a formal basis has also been expressed explicitly on occasion¹⁰. Thus, Boolos, Burgess and Jeffrey¹¹ define a diagonal function, d , any value of which can be decided effectively, although there is no single algorithm that can effectively compute d .

Now, the straightforward way of expressing this phenomenon should be to say that there are well-defined number-theoretic functions that are effectively computable instantiationally but not algorithmically. Yet, following Church and Turing, such functions are labeled as uncomputable¹²!

“According to Turing’s Thesis, since d is not Turing-computable, d cannot be effectively computable. Why not? After all, although no Turing machine computes the function d , we were able to compute at least its first few values. For since, as we have noted, $f_1 = f_1 = f_1 =$ the empty function we have $d(1) = d(2) = d(3) = 1$. And it may seem that we can actually compute $d(n)$ for any positive integer n —if we don’t run out of time.”¹³

The reluctance to treat a function such as $d(n)$ —or the function $\Omega(n)$ that computes the n^{th} digit in the decimal expression of a Chaitin constant Ω ¹⁴—as computable, on the grounds that the ‘time’ needed to compute it increases monotonically with n , is curious¹⁵; the same applies to any total Turing-computable function $f(n)$ ¹⁶!

1.2 Distinguishing between algorithmically verifiability and algorithmic computability

We now show in Theorem 1 that if Aristotle’s particularisation¹⁷ is presumed valid over the structure $\mathbb{N}$ of the natural numbers—as is the case under the standard interpretation of the first-order Peano Arithmetic PA—then it follows from the instantiational nature of the (constructively defined¹⁸) Gödel β -function that a primitive recursive relation can be instantiationally equivalent to an arithmetical relation, where the former is algorithmically computable over $\mathbb{N}$, whilst the latter is algorithmically verifiable but not algorithmically computable over $\mathbb{N}$.

¹⁰Parikh’s distinction between ‘decidability’ and ‘feasibility’ in [Pa71] also appears to echo the need for such a distinction.

¹¹[BBJ03], p. 37.

¹²The issue here seems to be that, when using language to express the abstract objects of our individual, and common, mental ‘concept spaces’, we use the word ‘exists’ loosely in three senses, without making explicit distinctions between them (see [An07c]).

¹³[BBJ03], p.37.

¹⁴Chaitin’s Halting Probability is given by $0 < \Omega = \sum 2^{-|p|} < 1$, where the summation is over all self-delimiting programs p that halt, and $|p|$ is the size in bits of the halting program p ; see [Ct75].

¹⁵The incongruity of this is addressed by Parikh in [Pa71].

¹⁶The only difference being that, in the latter case, we know there is a common ‘program’ of constant length that will compute $f(n)$ for any given natural number n ; in the former, we know we may need distinctly different programs for computing $f(n)$ for different values of n , where the length of the program will, sometime, reference n .

¹⁷See Definition 1 below.

¹⁸By Kurt Gödel; see Appendix A, Section 7, Lemma 1

Analogue distinctions in analysis: The distinction between algorithmically computable, and algorithmically verifiable but not algorithmically computable, number-theoretic functions seeks to reflect in arithmetic the essence of *uniform* methods¹⁹, classically characterised by the distinctions in analysis between: (a) uniformly continuous, and point-wise continuous but not uniformly continuous, functions over an interval; (b) uniformly convergent, and point-wise convergent but not uniformly convergent, series.

A limitation of set theory and a possible barrier to computation: We note, further, that the above distinction cannot be reflected within a language—such as the set theory ZF—which identifies ‘equality’ with ‘equivalence’. Since functions are defined extensionally as mappings, such a language cannot recognise that a set which represents a primitive recursive function may be equivalent to, but computationally different from, a set that represents an arithmetical function; where the former function is algorithmically computable over $\mathbb{N}$, whilst the latter is algorithmically verifiable but not algorithmically computable over $\mathbb{N}$.

1.2.1 Significance of Gödel’s β -function

In Theorem VII²⁰ of his seminal 1931 paper on formally undecidable arithmetical propositions, Gödel showed that, given a total number-theoretic function $f(x)$ and any natural number n , we can construct a primitive recursive function $\beta(z, y, x)$ and natural numbers b_n, c_n such that $\beta(b_n, c_n, i) = f(i)$ for all $0 \leq i \leq n$.

In this paper we shall essentially answer the following question affirmatively:

Query 1 *Does Gödel’s Theorem VII admit construction of an arithmetical function $A(x)$ such that:*

(a) *for any given natural number n , there is an algorithm that can verify $A(i) = f(i)$ for all $0 \leq i \leq n$ (hence $A(x)$ may be said to be algorithmically verifiable if $f(x)$ is recursive);*

(b) *there is no algorithm that can verify $A(i) = f(i)$ for all $0 \leq i$ (so $A(x)$ may be said to be algorithmically uncomputable)?*

1.2.2 Defining effective computability

We shall then formally define what it means for a formula of an arithmetical language to be:

- (i) Algorithmically verifiable;
- (ii) Algorithmically computable.

under an interpretation.

We shall further propose the definition:

Effective computability: A number-theoretic formula is effectively computable if, and only if, it is algorithmically verifiable.

¹⁹See Section 1.1.

²⁰[Go31], pp.30-31; reproduced below in Appendix A, Section 7.

Intuitionistically unobjectionable: We note first that since every finite set of integers is recursive, every well-defined number-theoretical formula is algorithmically verifiable, and so the above definition is intuitionistically unobjectionable; and second that the existence of an arithmetic formula that is algorithmically verifiable but not algorithmically computable (Theorem 1) supports Gödel’s reservations on Alonzo Church’s original intention to label his Thesis as a definition²¹.

We shall then show that the algorithmically verifiable and the algorithmically computable PA formulas are well-defined under the standard interpretation of PA since:

- (a) The PA-formulas are decidable as satisfied / unsatisfied or true / false under the standard interpretation of PA if, and only if, they are algorithmically verifiable;
- (b) The algorithmically computable PA-formulas are a proper subset of the algorithmically verifiable PA-formulas;
- (c) The PA-axioms are algorithmically computable as satisfied / true under the standard interpretation of PA;
- (d) Generalisation and Modus Ponens preserve algorithmically computable truth under the standard interpretation of PA;
- (e) The provable PA-formulas are precisely the ones that are algorithmically computable as satisfied / true under the standard interpretation of PA.

2 Comments, Notation and Standard Definitions

Comments We have taken some liberty in emphasising standard definitions selectively, and interspersing our arguments liberally with comments and references, generally of a foundational nature. These are intended to reflect our underlying thesis that essentially arithmetical problems appear more natural when expressed—and viewed—within the perspective of an interpretation of PA that appeals to the *evidence* provided by a deterministic Turing machine²² along the lines suggested in Section 6; a perspective that, by its very nature, cannot appeal implicitly to transfinite concepts.

Evidence “It is by now folklore ... that one can view the *values* of a simple functional language as specifying *evidence* for propositions in a constructive logic ...”²³.

Notation We use square brackets to indicate that the contents represent a symbol or a formula—of a formal theory—generally assumed to be well-formed unless otherwise indicated by the context.²⁴ We use an asterisk to indicate that

²¹See Section 1.1.

²²A deterministic Turing machine has only one possible move from a given configuration.

²³[Mu91].

²⁴In other words, expressions inside the square brackets are to be only viewed syntactically as juxtaposition of symbols that are to be formed and manipulated upon strictly in accordance with specific rules for such formation and manipulation—in the manner of a mechanical or electronic device—without any regards to what the symbolism might represent semantically under an interpretation that gives them meaning.

the expression is to be interpreted semantically with respect to some well-defined interpretation.

Definition 1 Aristotle’s particularisation *This holds that from an assertion such as:*

‘It is not the case that: for any given x , $P^*(x)$ does not hold’,

usually denoted symbolically by ‘ $\neg(\forall x)\neg P^*(x)$ ’, we may always validly infer in the classical, Aristotelean, logic of predicates²⁵ that:

‘There exists an unspecified x such that $P^*(x)$ holds’,

usually denoted symbolically by ‘ $(\exists x)P^*(x)$ ’.

The significance of Aristotle’s particularisation for the first-order predicate calculus: We note that in a formal language the formula ‘ $[(\exists x)P(x)]$ ’ is an abbreviation for the formula ‘ $[\neg(\forall x)\neg P(x)]$ ’. The commonly accepted interpretation of this formula—and a fundamental tenet of classical logic unrestrictedly adopted as intuitively obvious by standard literature that seeks to build upon the formal first-order predicate calculus²⁶—tacitly appeals to Aristotelean particularisation.

However, L. E. J. Brouwer had noted in his seminal 1908 paper on the unreliability of logical principles²⁷ that the commonly accepted interpretation of this formula is ambiguous if interpretation is intended over an infinite domain. He essentially argued that, even supposing the formula ‘ $[P(x)]$ ’ of a formal Arithmetical language interprets as an arithmetical relation denoted by ‘ $P^*(x)$ ’, and the formula ‘ $[\neg(\forall x)\neg P(x)]$ ’ as the arithmetical proposition denoted by ‘ $\neg(\forall x)\neg P^*(x)$ ’, the formula ‘ $[(\exists x)P(x)]$ ’ need not interpret as the arithmetical proposition denoted by the usual abbreviation ‘ $(\exists x)P^*(x)$ ’; and that such postulation is invalid as a general logical principle in the absence of a means for constructing some putative object a for which the proposition $P^*(a)$ holds in the domain of the interpretation.

Hence we shall follow the convention that the assumption that ‘ $(\exists x)P^*(x)$ ’ is the intended interpretation of the formula ‘ $[(\exists x)P(x)]$ ’—which is essentially the assumption that Aristotle’s particularisation holds over the domain of the interpretation—must always be explicit.

The significance of Aristotle’s particularisation for PA: In order to avoid intuitionistic objections to his reasoning, Gödel introduced the syntactic property of ω -consistency as an explicit assumption in his formal reasoning in his seminal 1931 paper on formally undecidable arithmetical propositions²⁸.

Gödel explained at some length²⁹ that his reasons for introducing ω -consistency explicitly was to avoid appealing to the semantic concept of classical arithmetical truth in Aristotle’s logic of predicates (which presumes Aristotle’s particularisation).

However, we note that the two concepts are meta-mathematically equivalent in the sense that, if PA is consistent, then PA is ω -consistent if, and only if, Aristotle’s particularisation holds under the standard interpretation of PA.

²⁵[HA28], pp.58-59.

²⁶See [Hi25], p.382; [HA28], p.48; [Sk28], p.515; [Go31], p.32.; [Kl52], p.169; [Ro53], p.90; [BF58], p.46; [Be59], pp.178 & 218; [Su60], p.3; [Wa63], p.314-315; [Qu63], pp.12-13; [Kn63], p.60; [Co66], p.4; [Me64], p.52(ii); [Nv64], p.92; [Li64], p.33; [Sh67], p.13; [Da82], p.xxv; [Rg87], p.xvii; [EC89], p.174; [Mu91]; [Sm92], p.18, Ex.3; [BBJ03], p.102.

²⁷[Br08].

²⁸[Go31], p.23 and p.28.

²⁹In his introduction on p.9 of [Go31].

Definition 2 *The structure $\mathbb{N}$* The structure of the natural numbers—namely, $\{N$ (the set of natural numbers); $=$ (equality); $'$ (the successor function); $+$ (the addition function); $*$ (the product function); 0 (the null element) $\}$.

Definition 3 *The axioms of first-order Peano Arithmetic (PA)*

$\text{PA}_1 [(x_1 = x_2) \rightarrow ((x_1 = x_3) \rightarrow (x_2 = x_3))];$
 $\text{PA}_2 [(x_1 = x_2) \rightarrow (x'_1 = x'_2)];$
 $\text{PA}_3 [0 \neq x'_1];$
 $\text{PA}_4 [(x'_1 = x'_2) \rightarrow (x_1 = x_2)];$
 $\text{PA}_5 [(x_1 + 0) = x_1];$
 $\text{PA}_6 [(x_1 + x'_2) = (x_1 + x_2)'];$
 $\text{PA}_7 [(x_1 \star 0) = 0];$
 $\text{PA}_8 [(x_1 \star x'_2) = ((x_1 \star x_2) + x_1)];$
 PA_9 For any well-formed formula $[F(x)]$ of PA:
 $[F(0) \rightarrow ((\forall x)(F(x) \rightarrow F(x'))) \rightarrow (\forall x)F(x)].$

Definition 4 *Generalisation in PA* If $[A]$ is PA-provable, then so is $[(\forall x)A]$.

Definition 5 *Modus Ponens in PA* If $[A]$ and $[A \rightarrow B]$ are PA-provable, then so is $[B]$.

Definition 6 *Standard interpretation of PA* The standard interpretation of PA over the structure $\mathbb{N}$ is the one in which the logical constants have their ‘usual’ interpretations³⁰ in Aristotle’s logic of predicates (which subsumes Aristotle’s particularisation), and³¹:

- (a) the set of non-negative integers is the domain;
- (b) the symbol $[0]$ interprets as the integer 0;
- (c) the symbol $[']$ interprets as the successor operation (addition of 1);
- (d) the symbols $[+]$ and $[*]$ interpret as ordinary addition and multiplication;
- (e) the symbol $[=]$ interprets as the identity relation.

3 Gödel’s Theorem VII and algorithmically verifiable, but not algorithmically computable, arithmetical propositions

In his seminal 1931 paper on formally undecidable arithmetical propositions, Gödel defined a curious primitive recursive function—Gödel’s β -function—as³²:

Definition 7 $\beta(x_1, x_2, x_3) = rm(1 + (x_3 + 1) \star x_2, x_1)$

where $rm(x_1, x_2)$ denotes the remainder obtained on dividing x_2 by x_1 .

Gödel showed that the above function has the remarkable property that:

Lemma 1 For any given denumerable sequence of natural numbers, say $f(k, 0), f(k, 1), \dots$, and any given natural number n , we can construct natural numbers b, c, j such that:

³⁰See [Me64], p.49.

³¹See [Me64], p.107.

³²cf. [Go31], p.31, Lemma 1; [Me64], p.131, Proposition 3.21.

- (i) $j = \max(n, f(k, 0), f(k, 1), \dots, f(k, n))$;
- (ii) $c = j!$;
- (iii) $\beta(b, c, i) = f(k, i)$ for $0 \leq i \leq n$.

Proof This is a standard result³³. □

Gödel's original argument: We reproduce Gödel's original argument—which yields this critical lemma—in Appendix A, Section 7.

Now we have the standard definition³⁴:

Definition 8 *A number-theoretic function $f(x_1, \dots, x_n)$ is said to be representable in PA if, and only if, there is a PA formula $[F(x_1, \dots, x_{n+1})]$ with the free variables $[x_1, \dots, x_{n+1}]$, such that, for any given natural numbers $k_1, \dots, k_{n+1}$:*

- (i) *if $f(k_1, \dots, k_n) = k_{n+1}$ then PA proves: $[F(k_1, \dots, k_n, k_{n+1})]$;*
- (ii) *PA proves: $[(\exists_1 x_{n+1})F(k_1, \dots, k_n, x_{n+1})]$.*

The function $f(x_1, \dots, x_n)$ is said to be strongly representable in PA if we further have that:

- (iii) *PA proves: $[(\exists_1 x_{n+1})F(x_1, \dots, x_n, x_{n+1})]$*

Interpretation of $[\exists_1]$: The symbol $[\exists_1]$ denotes ‘uniqueness’ under an interpretation which assumes that Aristotle’s particularisation holds in the domain of the interpretation. Formally, however, the PA formula $[(\exists_1 x_3)F(x_1, x_2, x_3)]$ is merely a short-hand notation for the PA formula $[\neg(\forall x_3)\neg F(x_1, x_2, x_3) \wedge (\forall y)(\forall z) (F(x_1, x_2, y) \wedge F(x_1, x_2, z) \rightarrow y = z)]$.

We then have:

Lemma 2 $\beta(x_1, x_2, x_3)$ is strongly represented in PA by $[Bt(x_1, x_2, x_3, x_4)]$, which is defined as follows:

$$[(\exists w)(x_1 = ((1 + (x_3 + 1) \star x_2) \star w + x_4) \wedge (x_4 < 1 + (x_3 + 1) \star x_2))].$$

Proof This is a standard result³⁵. □

Gödel further showed (also under the tacit, but critical, presumption of Aristotle’s particularisation³⁶) that:

Lemma 3 *If $f(x_1, x_2)$ is a recursive function defined by:*

- (i) $f(x_1, 0) = g(x_1)$
- (ii) $f(x_1, (x_2 + 1)) = h(x_1, x_2, f(x_1, x_2))$

³³cf. [Go31], p.31, p.31, Lemma 1; [Me64], p.131, Proposition 3.22.

³⁴[Me64], p.118.

³⁵cf. [Me64], p.131, proposition 3.21.

³⁶The implicit assumption being that the negation of a universally quantified formula of the first-order predicate calculus is indicative of “the existence of a counter-example”—[Go31], p.32.

where $g(x_1)$ and $h(x_1, x_2, x_3)$ are recursive functions of lower rank³⁷ that are represented in PA by well-formed formulas $[G(x_1, x_2)]$ and $[H(x_1, x_2, x_3, x_4)]$, then $f(x_1, x_2)$ is represented in PA by the following well-formed formula, denoted by $[F(x_1, x_2, x_3)]$:

$$[(\exists u)(\exists v)((\exists w)(Bt(u, v, 0, w) \wedge G(x_1, w)) \wedge Bt(u, v, x_2, x_3) \wedge (\forall w)(w < x_2 \rightarrow (\exists y)(\exists z)(Bt(u, v, w, y) \wedge Bt(u, v, (w+1), z) \wedge H(x_1, w, y, z))))].$$

Proof This is a standard result³⁸. □

Gödel's original argument: We reproduce Gödel's original argument and proof of this critical lemma in Appendix A, Section 7.

3.1 What does “ $[(\exists_1 x_3)F(k, m, x_3)]$ is provable” assert under the standard interpretation of PA?

Now, if the PA formula $[F(x_1, x_2, x_3)]$ represents in PA the recursive function denoted by $f(x_1, x_2)$ then by definition, for any given numerals $[k], [m]$, the formula $[(\exists_1 x_3)F(k, m, x_3)]$ is provable in PA; and true under the standard interpretation of PA. We thus have that:

Lemma 4 “ $[(\exists_1 x_3)F(k, m, x_3)]$ is true under the standard interpretation of PA” is the assertion that:

Given any natural numbers k, m , we can construct natural numbers $t_{(k,m)}, u_{(k,m)}, v_{(k,m)}$ —all functions of k, m —such that:

- (a) $\beta(u_{(k,m)}, v_{(k,m)}, 0) = g(k)$;
- (b) for all $i < m$, $\beta(u_{(k,m)}, v_{(k,m)}, i) = h(k, i, f(k, i))$;
- (c) $\beta(u_{(k,m)}, v_{(k,m)}, m) = t_{(k,m)}$;

where $f(x_1, x_2)$, $g(x_1)$ and $h(x_1, x_2, x_3)$ are any recursive functions that are formally represented in PA by $F(x_1, x_2, x_3)$, $G(x_1, x_2)$ and $H(x_1, x_2, x_3, x_4)$ respectively such that:

- (i) $f(k, 0) = g(k)$
- (ii) $f(k, (y + 1)) = h(k, y, f(k, y))$ for all $y < m$
- (iii) $g(x_1)$ and $h(x_1, x_2, x_3)$ are recursive functions that are assumed to be of lower rank than $f(x_1, x_2)$.

Proof For any given natural numbers k and m , if $[F(x_1, x_2, x_3)]$ interprets as a well-defined arithmetical relation under the standard interpretation of PA, then we can define a deterministic Turing machine TM that can ‘construct’ the sequences $f(k, 0), f(k, 1), \dots, f(k, m)$ and $\beta(u_{(k,m)}, v_{(k,m)}, 0), \beta(u_{(k,m)}, v_{(k,m)}, 1), \dots, \beta(u_{(k,m)}, v_{(k,m)}, m)$ and give evidence to verify the assertion. □

³⁷cf. [Me64], p.132; [Go31], p.30(2).

³⁸cf. [Go31], p.31(2); [Me64], p.132.

Does $[F(x_1, x_2, x_3)]$ interpret as a well-defined predicate? A critical issue that we do not address here is whether the PA formula $[F(x_1, x_2, x_3)]$ can be considered to interpret under a sound interpretation of PA as a well-defined predicate, since the denumerable sequences $\{f(k, 0), f(k, 1), \dots, f(k, m), m_p : p > 0 \text{ and } m_p \text{ is not equal to } m_q \text{ if } p \text{ is not equal to } q\}$ —are represented by denumerable, distinctly different, functions $\beta(u_{p_1}, v_{p_2}, i)$ respectively. There are thus denumerable pairs (u_{p_1}, v_{p_2}) for which $\beta(u_{p_1}, v_{p_2}, i)$ yields the sequence $f(k, 0), f(k, 1), \dots, f(k, m)$.

We now see that:

Theorem 1 *Under the standard interpretation of PA $[(\exists_1 x_3)F(x_1, x_2, x_3)]$ is algorithmically verifiable, but not algorithmically computable, as always true over $\mathbb{N}$.*

Proof It follows from Lemma 4 that:

(1) $[(\exists_1 x_3)F(k, m, x_3)]$ is PA-provable for any given numerals $[k, m]$. Hence $[(\exists_1 x_3)F(k, m, x_3)]$ is true under the standard interpretation of PA. It then follows from the definition of $[F(x_1, x_2, x_3)]$ in Lemma 3 that, for any given natural numbers k, m , we can construct some pair of natural numbers $u_{(k, m)}, v_{(k, m)}$ —where $u_{(k, m)}, v_{(k, m)}$ are functions of the given natural numbers k and m —such that:

- (a) $\beta(u_{(k, m)}, v_{(k, m)}, i) = f(k, i)$ for $0 \leq i \leq m$;
- (b) $F^*(k, m, f(k, m))$ holds in $\mathbb{N}$.

Since $\beta(x_1, x_2, x_3)$ is primitive recursive, $\beta(u_{(k, m)}, v_{(k, m)}, i)$ defines a deterministic Turing machine TM that can ‘construct’ the denumerable sequence $f'(k, 0), f'(k, 1), \dots$ for any given natural numbers k and m such that:

- (c) $f(k, i) = f'(k, i)$ for $0 \leq i \leq m$.

We can thus define a deterministic Turing machine TM that will give evidence that the PA formula $[(\exists_1 x_3)F(k, m, x_3)]$ is true under the standard interpretation of PA.

Hence $[(\exists_1 x_3)F(x_1, x_2, x_3)]$ is algorithmically verifiable over $\mathbb{N}$ under the standard interpretation of PA.

(2) Now, the pair of natural numbers $u_{(x_1, x_2)}, v_{(x_1, x_2)}$ are defined such that:

- (a) $\beta(u_{(x_1, x_2)}, v_{(x_1, x_2)}, i) = f(x_1, i)$ for $0 \leq i \leq x_2$;
- (b) $F^*(x_1, x_2, f(x_1, x_2))$ holds in $\mathbb{N}$;

where $v_{(x_1, x_2)}$ is defined in Lemma 3 as $j!$, and:

- (c) $j = \max(n, f(x_1, 0), f(x_1, 1), \dots, f(x_1, x_2))$;
- (d) n is the ‘number’ of terms in the sequence $f(x_1, 0), f(x_1, 1), \dots, f(x_1, x_2)$.

Since j is not definable for a denumerable sequence $\beta(u_{(x_1, x_2)}, v_{(x_1, x_2)})$,
i) we cannot define a denumerable sequence $f'(x_1, 0), f'(x_1, 1), \dots$
such that:

- (e) $f(k, i) = f'(k, i)$ for all $i \geq 0$.

We cannot thus define a deterministic Turing machine TM that will give evidence that the PA formula $[(\exists_1 x_3)F(x_1, x_2, x_3)]$ interprets as true under the standard interpretation of PA for any given sequence of numerals $[(a_1, a_2)]$.

Hence $[(\exists_1 x_3)F(x_1, x_2, x_3)]$ is not algorithmically computable over $\mathbb{N}$ under the standard interpretation of PA.

The theorem follows. □

The above theorem now suggests the following definition:

Definition 9 *Effective computability:* A number-theoretic function is *effectively computable* if, and only if, it is *algorithmically verifiable*.

However, we still need to formally define what it means for a number-theoretic formula of an arithmetical language to be:

- (i) Algorithmically verifiable;
- (ii) Algorithmically computable.

under an interpretation.

4 Interpretation of an arithmetical language in terms of Turing computability

We begin by noting that we can, in principle, define the classical ‘satisfaction’ and ‘truth’ of the formulas of a first order arithmetical language, such as PA, *verifiably* under an interpretation using as *evidence* the computations of a simple functional language.

Such definitions follow straightforwardly for the atomic formulas of the language (i.e., those without the logical constants that correspond to ‘negation’, ‘conjunction’, ‘implication’ and ‘quantification’) from the standard definition of a deterministic Turing machine³⁹, based essentially on Alan Turing’s seminal 1936 paper on computable numbers⁴⁰.

Moreover, it follows from Alfred Tarski’s seminal 1933 paper on the the concept of truth in the languages of the deductive sciences⁴¹ that the ‘satisfaction’

³⁹[Me64], pp.229-231.

⁴⁰[Tu36].

⁴¹[Ta33].

and ‘truth’ of those formulas of a first-order language which contain logical constants can be inductively defined, under an interpretation, in terms of the ‘satisfaction’ and ‘truth’ of the interpretations of only the atomic formulas of the language.

Hence the ‘satisfaction’ and ‘truth’ of those formulas (of an arithmetical language) which contain logical constants can, in principle, also be defined verifiably under an interpretation using as evidence the computations of a deterministic Turing machine.

We show in Section 6 that this is indeed the case for PA under the standard interpretation $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$, when this is explicitly defined as in Section 6.1.

We show in Section 6.2, moreover, that we can further define ‘algorithmic truth’ and ‘algorithmic falsehood’ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ such that the PA axioms interpret as always algorithmically true.

Significance of ‘algorithmic truth’: The *algorithmically* true propositions of $\mathbb{N}$ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ are thus a proper subset of the *verifiably* true propositions of $\mathbb{N}$ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$; and suggest a possible finitary ‘model’⁴² of PA that would establish the consistency of PA constructively.

4.1 The definitions of ‘algorithmic truth’ and ‘algorithmic falsehood’ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ are not symmetric with respect to ‘truth’ and ‘falsehood’ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$

However, the definitions (in Section 6.2) of ‘algorithmic truth’ and ‘algorithmic falsehood’ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ are not symmetric with respect to classical (verifiable) ‘truth’ and ‘falsehood’ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.

For instance, if a formula $[F(x_1, x_2, \dots, x_n)]$ of an arithmetic is algorithmically true under an interpretation (such as $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$) that appeals to the evidence provided by the computations of a deterministic Turing machine, then, for any given denumerable sequence of numerical values $[a_1, a_2, \dots]$, the formula $[F(a_1, a_2, \dots, a_n)]$ is also algorithmically true under the interpretation.

Denumerable sequence: We shall presume that any such sequence is ‘given’ in the sense of being defined by the ‘evidence’ of a deterministic Turing machine.

In other words, there is a deterministic Turing machine which can provide evidence that the interpretation $F^*(a_1, a_2, \dots, a_n)$ of $[F(a_1, a_2, \dots, a_n)]$ *holds* in $\mathbb{N}$ for any given denumerable sequence of natural numbers $(a_1, a_2, \dots)$.

Defining the term ‘hold’: We define the term ‘hold’—when used in connection with an interpretation of a formal language and, more specifically, with reference to the operations of a deterministic Turing machine associated with the atomic formulas of the language—explicitly in Section 6; the aim being to avoid appealing to the classically subjective (and existential) connotation implicitly associated with the term under an implicitly defined standard interpretation of an arithmetic⁴³.

However, if a formula $[F(x_1, x_2, \dots, x_n)]$ of an arithmetic is algorithmically false under an interpretation that appeals to the evidence provided by the computations of a deterministic Turing machine, we cannot conclude that there is

⁴²[Me64], p.51.

⁴³As, for instance, in [Go31].

a denumerable sequence of numerical values $[a_1, a_2, \dots]$ such that the formula $[F(a_1, a_2, \dots, a_n)]$ is algorithmically false under the interpretation.

Reason: If a formula $[F(x_1, x_2, \dots, x_n)]$ of an arithmetic is algorithmically false under such an interpretation, then we can only conclude that there is no deterministic Turing machine which can provide evidence that the interpretation $F^*(x_1, x_2, \dots, x_n)$ holds in $\mathbb{N}$ for any given denumerable sequence of natural numbers $(a_1, a_2, \dots)$; we cannot conclude that there is a denumerable sequence of natural numbers $(b_1, b_2, \dots)$ such that $F^*(b_1, b_2, \dots, b_n)$ does not hold in $\mathbb{N}$.

Such a conclusion would require:

(i) either some additional evidence that will verify for some assignment of numerical values to the free variables of $[F]$ that the corresponding interpretation F^* does not hold⁴⁴;

(ii) or the additional assumption that either Aristotle's particularisation holds over the domain of the interpretation (as is implicitly presumed under the standard interpretation of PA) or the arithmetic is ω -consistent⁴⁵.

An issue of consistency: An issue that we do not address here is whether the assumption of Aristotle's particularisation (or that of ω -completeness) is consistent with the evidence provided by the computations of a deterministic Turing machine for defining the satisfaction and truth of the formulas of an arithmetic under an interpretation.

5 Defining algorithmic verifiability and algorithmic computability

The asymmetry of Section 4.1 suggests the following two concepts⁴⁶:

Definition 10 *Algorithmic verifiability:* *An arithmetical formula $[F(x_1, x_2, \dots, x_n)]$ is algorithmically verifiable under an interpretation if, and only if, for any given sequence of numerals $[a_1, a_2, \dots, a_n]$, we can define a deterministic Turing machine TM that computes $[F(a_1, a_2, \dots, a_n)]$ and will halt on null input if, and only if, $[F(a_1, a_2, \dots, a_n)]$ interprets as either true or false under the interpretation.*

Interpretation of an arithmetical language *verifiably* in terms of Turing computability Of course such a definition requires defining the 'algorithmic verifiability' of the formulas of an arithmetical language under an interpretation in terms of the operations of a deterministic Turing machine. Such definitions follow straightforwardly for the atomic formulas of the language (i.e., those without the logical constants that correspond to 'negation', 'conjunction', 'implication' and 'quantification') from Alan Turing's seminal 1936 paper on computable numbers⁴⁷. Moreover, it follows from Alfred Tarski's seminal 1933 paper on the concept of truth in the languages of the deductive sciences⁴⁸ that the 'algorithmic verifiability' of the formulas of a formal language which contain logical constants can be inductively defined under an interpretation in terms of the 'algorithmic verifiability' of the interpretations of the atomic formulas of the language. Hence the 'algorithmic verifiability' of the formulas containing logical constants in the above definition can, in principle, be defined in terms of the operations of a deterministic Turing machine. We show in Section 6 that this is indeed the case.

⁴⁴Essentially reflecting Brouwer's objection to the assumption of Aristotle's particularisation over an infinite domain.

⁴⁵An assumption explicitly introduced by Gödel in [Go31].

⁴⁶My thanks to Dr. Chaitanya H. Mehta for advising that the focus of this investigation should be the distinction between these two concepts.

⁴⁷[Tu36].

⁴⁸[Ta33].

Definition 11 *Algorithmic computability:* *An arithmetical formula $[F(x_1, x_2, \dots, x_n)]$ is algorithmically computable under an interpretation if, and only if, we can define a deterministic Turing machine TM that computes $[F(x_1, x_2, \dots, x_n)]$ and, for any given sequence of numerals $[a_1, a_2, \dots, a_n]$, will halt on $[a_1, a_2, \dots, a_n]$ if, and only if, $[F(a_1, a_2, \dots, a_n)]$ interprets as either true or false under the interpretation.*

Interpretation of an arithmetical language algorithmically in terms of Turing computability We show in Section 6 that the ‘algorithmic computability’ of the formulas of a formal language which contain logical constants can also be inductively defined under an interpretation in terms of the ‘algorithmic computability’ of the interpretations of the atomic formulas of the language.

We now show that the above concepts are well-defined under the standard interpretation of PA.

6 Standard definitions of the ‘satisfiability’ and ‘truth’ of a formal language under an interpretation

We note that standard interpretations of the formal reasoning and conclusions of classical first order theory—based primarily on the work of Cantor, Gödel, Tarski, and Turing—seem to admit the impression that the classical, Tarskian, truth (satisfiability) of the propositions of a formal mathematical language under an interpretation is, both, non-algorithmic and essentially unverifiable constructively⁴⁹.

However—if mathematics is to serve as a universal set of languages of, both, precise expression and unambiguous communication—such interpretations may need to be balanced by an alternative, constructive and intuitionistically unobjectionable, interpretation—of classical foundational concepts—in which non-algorithmic truth (satisfiability) is defined effectively⁵⁰.

For instance, we note that—essentially following standard expositions⁵¹ of Tarski’s inductive definitions on the ‘satisfiability’ and ‘truth’ of the formulas of a formal language under an interpretation—we can define:

Definition 12 *If $[A]$ is an atomic formula $[A(x_1, x_2, \dots, x_n)]$ of a formal language S , then the denumerable sequence $(a_1, a_2, \dots)$ in the domain $\mathbb{D}$ of an interpretation $\mathcal{I}_{S(\mathbb{D})}$ of S satisfies $[A]$ if, and only if:*

(i) *$[A(x_1, x_2, \dots, x_n)]$ interprets under $\mathcal{I}_{S(\mathbb{D})}$ as a relation $A^*(x_1, x_2, \dots, x_n)$ in $\mathbb{D}$ for a witness $\mathcal{W}_{\mathbb{D}}$ of $\mathbb{D}$;*

(ii) *there is a Satisfaction Method, $SM(\mathcal{I}_{S(\mathbb{D})})$ that provides objective evidence⁵² by which any witness $\mathcal{W}_{\mathbb{D}}$ of $\mathbb{D}$ can **define** for any atomic formula $[A(x_1, x_2, \dots, x_n)]$ of S , and any given denumerable sequence $(b_1, b_2, \dots)$ of $\mathbb{D}$, whether the proposition $A^*(b_1, b_2, \dots, b_n)$ holds or not in $\mathbb{D}$;*

⁴⁹Thus giving rise to J. R. Lucas’ Gödelian Argument in [Lu61]; but see also [An07a], [An07b], [An07c] and [An08].

⁵⁰Such an interpretation is broadly outlined in [An07d]

⁵¹cf. [Me64], p.51.

⁵²In the sense of [Mu91].

(iii) $A^*(a_1, a_2, \dots, a_n)$ holds in $\mathbb{D}$ for $\mathcal{W}_{\mathbb{D}}$.

Witness: From a constructive perspective, the existence of a ‘witness’ as in (i) above is implicit in the usual expositions of Tarski’s definitions.

Satisfaction Method: From a constructive perspective, the existence of a Satisfaction Method as in (ii) above is also implicit in the usual expositions of Tarski’s definitions.

A constructive perspective: We highlight the word ‘*define*’ in (ii) above to emphasise the constructive perspective underlying this paper; which is that the concepts of ‘satisfaction’ and ‘truth’ under an interpretation are to be explicitly viewed as objective assignments by a convention that is witness-independent. A Platonist perspective would substitute ‘decide’ for ‘define’, thus implicitly suggesting that these concepts can ‘exist’, in the sense of needing to be discovered by some witness-dependent means—eerily akin to a ‘revelation’—even when the domain $\mathbb{D}$ is $\mathbb{N}$.

We can now inductively assign truth values of ‘satisfaction’, ‘truth’, and ‘falsity’ to the compound formulas of a first-order theory S under the interpretation $\mathcal{I}_{S(\mathbb{D})}$ in terms of *only* the satisfiability of the atomic formulas of S over $\mathbb{D}$ as follows⁵³:

Definition 13 A denumerable sequence s of $\mathbb{D}$ satisfies $[\neg A]$ under $\mathcal{I}_{S(\mathbb{D})}$ if, and only if, s does not satisfy $[A]$;

Definition 14 A denumerable sequence s of $\mathbb{D}$ satisfies $[A \rightarrow B]$ under $\mathcal{I}_{S(\mathbb{D})}$ if, and only if, either it is not the case that s satisfies $[A]$, or s satisfies $[B]$;

Definition 15 A denumerable sequence s of $\mathbb{D}$ satisfies $[(\forall x_i)A]$ under $\mathcal{I}_{S(\mathbb{D})}$ if, and only if, given any denumerable sequence t of $\mathbb{D}$ which differs from s in at most the i ’th component, t satisfies $[A]$;

Definition 16 A well-formed formula $[A]$ of $\mathbb{D}$ is true under $\mathcal{I}_{S(\mathbb{D})}$ if, and only if, given any denumerable sequence t of $\mathbb{D}$, t satisfies $[A]$;

Definition 17 A well-formed formula $[A]$ of $\mathbb{D}$ is false under $\mathcal{I}_{S(\mathbb{D})}$ if, and only if, it is not the case that $\mathbb{D}$ is true under $\mathcal{I}_{S(\mathbb{D})}$.

It follows that⁵⁴:

Theorem 2 (*Satisfaction Theorem*) If, for any interpretation $\mathcal{I}_{S(\mathbb{D})}$ of a first-order theory S , there is a Satisfaction Method $SM(\mathcal{I}_{S(\mathbb{D})})$ which holds for a witness $\mathcal{W}_{\mathbb{D}}$ of $\mathbb{D}$, then:

(i) The Δ_0 formulas of S are decidable as either true or false over $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$;

(ii) If the Δ_n formulas of S are decidable as either true or as false over $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$, then so are the $\Delta(n+1)$ formulas of S .

⁵³Compare [Me64], p.51; [Mu91].

⁵⁴cf. [Me64], pp.51–53.

Proof It follows from the above definitions that:

(a) If, for any given atomic formula $[A(x_1, x_2, \dots, x_n)]$ of S , it is decidable by $\mathcal{W}_{\mathbb{D}}$ whether or not a given denumerable sequence $(a_1, a_2, \dots)$ of $\mathbb{D}$ satisfies $[A(x_1, x_2, \dots, x_n)]$ in $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$ then, for any given compound formula $[A^1(x_1, x_2, \dots, x_n)]$ of S containing any one of the logical constants $\neg, \rightarrow, \forall$, it is decidable by $\mathcal{W}_{\mathbb{D}}$ whether or not $(a_1, a_2, \dots)$ satisfies $[A^1(x_1, x_2, \dots, x_n)]$ in $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$;

(b) If, for any given compound formula $[B^n(x_1, x_2, \dots, x_n)]$ of S containing n of the logical constants $\neg, \rightarrow, \forall$, it is decidable by $\mathcal{W}_{\mathbb{D}}$ whether or not a given denumerable sequence $(a_1, a_2, \dots)$ of $\mathbb{D}$ satisfies $[B^n(x_1, x_2, \dots, x_n)]$ in $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$ then, for any given compound formula $[B^{(n+1)}(x_1, x_2, \dots, x_n)]$ of S containing $n+1$ of the logical constants $\neg, \rightarrow, \forall$, it is decidable by $\mathcal{W}_{\mathbb{D}}$ whether or not $(a_1, a_2, \dots)$ satisfies $[B^{(n+1)}(x_1, x_2, \dots, x_n)]$ in $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$;

We thus have that:

(c) The Δ_0 formulas of S are decidable by $\mathcal{W}_{\mathbb{D}}$ as either true or false over $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$;

(d) If the Δ_n formulas of S are decidable by $\mathcal{W}_{\mathbb{D}}$ as either true or as false over $\mathbb{D}$ under $\mathcal{I}_{S(\mathbb{D})}$, then so are the $\Delta(n+1)$ formulas of S . $\square$

In other words, if the atomic formulas of S interpret under $\mathcal{I}_{S(\mathbb{D})}$ as decidable with respect to the Satisfaction Method $\text{SM}(\mathcal{I}_{S(\mathbb{D})})$ by a witness $\mathcal{W}_{\mathbb{D}}$ over some domain $\mathbb{D}$, then the propositions of S (i.e., the Π_n and Σ_n formulas of S) also interpret as decidable with respect to $\text{SM}(\mathcal{I}_{S(\mathbb{D})})$ by the witness $\mathcal{W}_{\mathbb{D}}$ over $\mathbb{D}$.

6.1 An explicit definition of the standard interpretation of PA

We now consider the application of Tarski's definitions to the standard interpretation $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ of PA where:

(a) we define S as PA with standard first-order predicate calculus as the underlying logic;

(b) we define $\mathbb{D}$ as $\mathbb{N}$;

(c) we take $\text{SM}(\mathcal{I}_{PA(\mathbb{N}, \text{Standard})})$ as a deterministic Turing machine.

We note that:

Theorem 3 *The atomic formulas of PA are algorithmically verifiable under the standard interpretation $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.*

Proof If $[A(x_1, x_2, \dots, x_n)]$ is an atomic formula of PA then, for any given denumerable sequence of numerals $[b_1, b_2, \dots]$, the PA formula $[A(b_1, b_2, \dots, b_n)]$ is an atomic formula of the form $[c = d]$, where $[c]$ and $[d]$ are atomic PA formulas that denote PA numerals. Since $[c]$ and $[d]$ are recursively defined formulas in the language of PA, it follows from a standard result⁵⁵ that, if PA is consistent,

⁵⁵For any natural numbers m, n , if $m \neq n$, then PA proves $[\neg(m = n)]$ ([Me64], p.110, Proposition 3.6). The converse is obviously true.

then $[c = d]$ interprets as the proposition $c = d$ which either holds or not for a witness $\mathcal{W}_{\mathbb{N}}$ in $\mathbb{N}$.

Hence, if PA is consistent, then $[A(x_1, x_2, \dots, x_n)]$ is algorithmically verifiable since, for any given denumerable sequence of numerals $[b_1, b_2, \dots]$, we can define a deterministic Turing machine TM that will compute $[A(b_1, b_2, \dots, b_n)]$ and halt on null input as evidence that the PA formula $[A(b_1, b_2, \dots, b_n)]$ is decidable under the interpretation.

The theorem follows. $\square$

We thus have that:

Corollary 1 *The ‘satisfaction’ and ‘truth’ of PA formulas containing logical constants can be defined under the standard interpretation of PA in terms of the evidence provided by the computations of a deterministic Turing machine.*

6.2 Defining ‘algorithmic truth’ under the standard interpretation of PA and interpreting the PA axioms

Now we note that, in addition to Theorem 3:

Theorem 4 *The atomic formulas of PA are algorithmically computable under the standard interpretation $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.*

Proof If $[A(x_1, x_2, \dots, x_n)]$ is an atomic formula of PA then we can define a deterministic Turing machine TM that will compute $[A(x_1, x_2, \dots, x_n)]$ and halt on any given denumerable sequence of numerals $[b_1, b_2, \dots]$ as evidence that the PA formula $[A(b_1, b_2, \dots, b_n)]$ is decidable under the interpretation.

The theorem follows. $\square$

This suggests the following definitions:

Definition 18 *A well-formed formula $[A]$ of PA is algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ if, and only if, there is a deterministic Turing machine which computes $[A]$ and provides evidence that, given any denumerable sequence t of $\mathbb{N}$, t satisfies $[A]$;*

Definition 19 *A well-formed formula $[A]$ of PA is algorithmically false under $\mathcal{I}_{PA(\mathbb{N})}$ if, and only if, it is not algorithmically true under $\mathcal{I}_{PA(\mathbb{N})}$.*

The significance of defining ‘algorithmic truth’ under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ as above is that:

Lemma 5 *The PA axioms PA_1 to PA_8 are algorithmically computable as algorithmically true over $\mathbb{N}$ under the interpretation $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.*

Proof Since $[x + y]$, $[x \star y]$, $[x = y]$, $[x']$ are defined recursively⁵⁶, the PA axioms PA_1 to PA_8 interpret as recursive relations that do not involve any quantification. The lemma follows straightforwardly from Definitions 12 to 17 in Section 6 and Theorem 3. $\square$

⁵⁶cf. [Go31], p.17.

Lemma 6 *For any given PA formula $[F(x)]$, the Induction axiom schema $[F(0) \rightarrow ((\forall x)(F(x) \rightarrow F(x')))) \rightarrow (\forall x)F(x)]$ interprets as algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.*

Proof By Definitions 12 to 19:

(a) If $[F(0)]$ interprets as algorithmically false under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ the lemma is proved.

Since $[F(0) \rightarrow ((\forall x)(F(x) \rightarrow F(x')))) \rightarrow (\forall x)F(x)]$ interprets as algorithmically true if, and only if, either $[F(0)]$ interprets as algorithmically false or $[((\forall x)(F(x) \rightarrow F(x')))) \rightarrow (\forall x)F(x)]$ interprets as algorithmically true.

(b) If $[F(0)]$ interprets as algorithmically true and $[((\forall x)(F(x) \rightarrow F(x')))) \rightarrow (\forall x)F(x)]$ interprets as algorithmically false under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$, the lemma is proved.

(c) If $[F(0)]$ and $[((\forall x)(F(x) \rightarrow F(x')))) \rightarrow (\forall x)F(x)]$ both interpret as algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$, then by Definition 18 there is a deterministic Turing machine TM that, for any natural number n , will give evidence that the formula $[F(n) \rightarrow F(n')]$ is true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.

Since $[F(0)]$ interprets as algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$, it follows that there is a deterministic Turing machine TM that, for any natural number n , will give evidence that the formula $[F(n)]$ is true under the interpretation.

Hence $[(\forall x)F(x)]$ is algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.

Since the above cases are exhaustive, the lemma follows. $\square$

The Poincaré-Hilbert debate: We note that Lemma 6 appears to settle the Poincaré-Hilbert debate⁵⁷ in the latter's favour. Poincaré believed that the Induction Axiom could not be justified finitarily, as any such argument would necessarily need to appeal to infinite induction. Hilbert believed that a finitary proof of the consistency of PA was possible.

Lemma 7 *Generalisation preserves algorithmic truth under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.*

Proof The two meta-assertions:

' $[F(x)]$ interprets as algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ '⁵⁸,

and

' $[(\forall x)F(x)]$ interprets as algorithmically true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$ '

both mean:

$[F(x)]$ is algorithmically computable as always true under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$. $\square$

⁵⁷See [Hi27], p.472; also [Br13], p.59; [We27], p.482; [Pa71], p.502-503.

⁵⁸See Definition 16

It is also straightforward to see that:

Lemma 8 *Modus Ponens preserves algorithmic truth under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$.*
□

We thus have that:

Theorem 5 *The axioms of PA are always algorithmically true under the interpretation $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$, and the rules of inference of PA preserve the properties of algorithmic satisfaction/truth under $\mathcal{I}_{PA(\mathbb{N}, \text{Standard})}$, without appeal to Aristotle's particularisation.* □

7 Appendix A: Gödel's Theorem VII(2)

(Excerpted from [Go31] pp.29-31.)

Every relation of the form $x_0 = \phi(x_1, \dots, x_n)$, where ϕ is recursive, is arithmetical and we apply complete induction on the rank of ϕ . Let ϕ have rank s ($s > 1$). ...

$$\begin{aligned}\phi(0, x_2, \dots, x_n) &= \psi(x_2, \dots, x_n) \\ \phi(k+1, x_2, \dots, x_n) &= \mu[k, \phi(k, x_2, \dots, x_n), x_2, \dots, x_n] \\ \text{(where } \psi, \mu \text{ have lower rank than } s\text{).}\end{aligned}$$

...we apply the following procedure: one can express the relation $x_0 = \phi(x_1, \dots, x_n)$ with the help of the concept "sequence of numbers" (f)⁵⁹ in the following manner:

$$\begin{aligned}x_0 = \phi(x_1, \dots, x_n) &\sim (\exists f)\{f_0 = \psi(x_2, \dots, x_n) \ \& \ (\forall k)(k < x_1 \rightarrow \\ f_{k+1} &= \mu(k, f_k, x_2, \dots, x_n) \ \& \ x_0 = f_{x_1})\}\end{aligned}$$

If $S(y, x_2, \dots, x_n), T(z, x_1, \dots, x_{n+1})$ are the arithmetical relations which, according to the inductive hypothesis, are equivalent to $y = \psi(x_2, \dots, x_n)$, and $z = \mu(x_1, \dots, x_{n+1})$ respectively, then we have:

$$\begin{aligned}x_0 = \phi(x_1, \dots, x_n) &\sim (\exists f)\{S(f_0, x_2, \dots, x_n) \ \& \ (\forall k)[k < x_1 \rightarrow \\ T(f_{k+1}, k, x_2, \dots, x_n)] &\ \& \ x_0 = f_{x_1}\}\end{aligned} \quad (17)$$

Now we replace the concept "sequence of numbers" by "pairs of numbers" by correlating with the number pair n, d the sequence of numbers $f^{(n,d)}$ ($f_k^{(n,d)} = [n]_{1+(k+1)d}$, where $[n]_p$ denotes the smallest non-negative remainder of n modulo p).

Then:

Lemma 1: If f is an arbitrary sequence of natural numbers and k is an arbitrary natural number, then there exists a pair of natural numbers n, d such that $f^{(n,d)}$ and f coincide in their first k terms.

Proof: Let l be the greatest of the numbers $k, f_0, f_1, \dots, f_{k-1}$. Determine n so that

⁵⁹ f denotes here a variable whose domain is the sequence of natural numbers. The $(k+1)$ st term of a sequence f is designated f_k (and the first, f_0).

$$n \equiv f_i \pmod{1 + (i+1)l!} \text{ for } i = 0, 1, \dots, k-1,$$

which is possible, since any two of the numbers $1 + (i+1)l!$ ($i = 0, 1, \dots, k-1$) are relatively prime. For, a prime dividing two of these numbers must also divide the difference $(i_1 - i_2)l!$ and therefore, since $i_1 - i_2 < l$, must also divide $l!$, which is impossible. The number pair $n, l!$ fulfills our requirement.

Since the relation $x = [n]_p$ is defined by

$$x \equiv n \pmod{p} \text{ \& } x < p$$

and is therefore arithmetical, then so also is the relation $P(x_0, x_1, \dots, x_n)$ defined as follows:

$$P(x_0, x_1, \dots, x_n) \equiv (\exists n, d) \{ S([n]_{d+1}, x_2, \dots, x_n) \text{ \& } (\forall k) [k < x_1 \rightarrow T([n]_{1+d(k+2)}, k, [n]_{1+d(k+1)}, x_2, \dots, x_n)] \text{ \& } x_0 = [n]_{1+d(x_1+1)} \}$$

which, according to (17) and Lemma 1, is equivalent to $x_0 = \phi(x_1, \dots, x_n)$ (in the sequence f in (17) only its values up to the $(x+1)$ th term matter). Thus, Theorem VII(2) is proved.

Comment: Gödel's remark that "in the sequence f in (17) only its values up to the $(x+1)$ th term matter" is significant. The proof of Theorem 1 depends upon the fact that the equivalence between $f^{(n,d)}$ and f cannot be extended non-terminatingly.

References

- [BBJ03] George S. Boolos, John P. Burgess, Richard C. Jeffrey. 2003. *Computability and Logic* (4th ed). Cambridge University Press, Cambridge.
- [Be59] Evert W. Beth. 1959. *The Foundations of Mathematics*. Studies in Logic and the Foundations of Mathematics. Edited by L. E. J. Brouwer, E. W. Beth, A. Heyting. 1959. North Holland Publishing Company, Amsterdam.
- [BF58] Paul Bernays and Abraham A. Fraenkel. 1958. *Axiomatic Set Theory*. Studies in Logic and the Foundations of Mathematics. Edited by L. E. J. Brouwer, E. W. Beth, A. Heyting. 1959. North Holland Publishing Company, Amsterdam.
- [Br08] L. E. J. Brouwer. 1908. *The Unreliability of the Logical Principles*. English translation in A. Heyting, Ed. L. E. J. Brouwer: *Collected Works 1: Philosophy and Foundations of Mathematics*. Amsterdam: North Holland / New York: American Elsevier (1975): pp. 107-111.
- [Br13] L. E. J. Brouwer. 1913. *Intuitionism and Formalism*. Inaugural address at the University of Amsterdam, October 14, 1912. Translated by Professor Arnold Dresden for the Bulletin of the American Mathematical Society, Volume 20 (1913), pp.81-96. 1999. Electronically published in Bulletin (New Series) of the American Mathematical Society, Volume 37, Number 1, pp.55-64.
- [Ch36] Alonzo Church. 1936. *An unsolvable problem of elementary number theory*. In M. Davis (ed.). 1965. *The Undecidable*. Raven Press, New York. Reprinted from the Am. J. Math., Vol. 58, pp.345-363.
- [Co66] Paul J. Cohen. 1966. *Set Theory and the Continuum Hypothesis*. (Lecture notes given at Harvard University, Spring 1965) W. A. Benjamin, Inc., New York.
- [Ct75] Gregory J. Chaitin. 1975. *A Theory of Program Size Formally Identical to Information Theory*. J. Assoc. Comput. Mach. 22 (1975), pp. 329-340.
- [Da82] Martin Davis. 1958. *Computability and Unsolvability*. 1982 ed. Dover Publications, Inc., New York.

- [EC89] Richard L. Epstein, Walter A. Carnielli. 1989. *Computability: Computable Functions, Logic, and the Foundations of Mathematics*. Wadsworth & Brooks, California.
- [Go31] Kurt Gödel. 1931. *On formally undecidable propositions of Principia Mathematica and related systems I*. Translated by Elliott Mendelson. In M. Davis (ed.). 1965. *The Undecidable*. Raven Press, New York.
- [Go51] Kurt Gödel. 1951. *Some basic theorems on the foundations of mathematics and their implications*. Gibbs lecture. In Kurt Gödel, Collected Works III, pp.304-323. 1995. *Unpublished Essays and Lectures*. Solomon Feferman et al (ed.). Oxford University Press, New York.
- [HA28] David Hilbert & Wilhelm Ackermann. 1928. *Principles of Mathematical Logic*. Translation of the second edition of the *Grundzüge Der Theoretischen Logik*. 1928. Springer, Berlin. 1950. Chelsea Publishing Company, New York.
- [Hi25] David Hilbert. 1925. *On the Infinite*. Text of an address delivered in Münster on 4th June 1925 at a meeting of the Westphalian Mathematical Society. In Jean van Heijenoort. 1967. Ed. *From Frege to Gödel: A source book in Mathematical Logic, 1878 - 1931*. Harvard University Press, Cambridge, Massachusetts.
- [Hi27] David Hilbert. 1927. *The Foundations of Mathematics*. Text of an address delivered in July 1927 at the Hamburg Mathematical Seminar. In Jean van Heijenoort. 1967. Ed. *From Frege to Gödel: A source book in Mathematical Logic, 1878 - 1931*. Harvard University Press, Cambridge, Massachusetts.
- [Kl52] Stephen Cole Kleene. 1952. *Introduction to Metamathematics*. North Holland Publishing Company, Amsterdam.
- [Kn63] G. T. Kneebone. 1963. *Mathematical Logic and the Foundations of Mathematics: An Introductory Survey*. D. Van Norstrand Company Limited, London.
- [Li64] A. H. Lightstone. 1964. *The Axiomatic Method*. Prentice Hall, NJ.
- [Lu61] J. R. Lucas. 1961. *Minds, Machines and Gödel*. Philosophy, XXXVI, 1961, pp.112-127; reprinted in *The Modeling of Mind*. Kenneth M.Sayre and Frederick J.Crosson, eds., Notre Dame Press, 1963, pp.269-270; and *Minds and Machines*, ed. Alan Ross Anderson, Prentice-Hall, 1954, pp.43-59.
- [Me64] Elliott Mendelson. 1964. *Introduction to Mathematical Logic*. Van Norstrand, Princeton.
- [Mu91] Chetan R. Murthy. 1991. *An Evaluation Semantics for Classical Proofs*. Proceedings of Sixth IEEE Symposium on Logic in Computer Science, pp. 96-109, (also Cornell TR 91-1213), 1991.
- [Nv64] P. S. Novikov. 1964. *Elements of Mathematical Logic*. Oliver & Boyd, Edinburgh and London.
- [Pa71] Rohit Parikh. 1971. *Existence and Feasibility in Arithmetic*. The Journal of Symbolic Logic, Vol.36, No. 3 (Sep., 1971), pp. 494-508.
- [Qu63] Willard Van Orman Quine. 1963. *Set Theory and its Logic*. Harvard University Press, Cambridge, Massachusette.
- [Rg87] Hartley Rogers Jr. 1987. *Theory of Recursive Functions and Effective Computability*. MIT Press, Cambridge, Massachusetts.
- [Ro36] J. Barkley Rosser. 1936. *Extensions of some Theorems of Gödel and Church*. In M. Davis (ed.). 1965. *The Undecidable*. Raven Press, New York. Reprinted from The Journal of Symbolic Logic. Vol.1. pp.87-91.
- [Ro53] J. Barkley Rosser. 1953. *Logic for Mathematicians*. McGraw Hill, New York.

- [Sh67] Joseph R. Shoenfield. 1967. *Mathematical Logic*. Reprinted 2001. A. K. Peters Ltd., Massachusetts.
- [Si97] Wilfried Sieg. 1997. *Step by recursive step: Church's analysis of effective calculability*. Bulletin of Symbolic Logic, Volume 3, Number 2.
- [Sk28] Thoralf Skolem. 1928. *On Mathematical Logic*. Text of a lecture delivered on 22nd October 1928 before the Norwegian Mathematical Association. In Jean van Heijenoort. 1967. Ed. *From Frege to Gödel: A source book in Mathematical Logic, 1878 - 1931*. Harvard University Press, Cambridge, Massachusetts.
- [Sm92] Raymond M. Smullyan. 1992. *Gödel's Incompleteness Theorems*. Oxford University Press, Inc., New York.
- [Su60] Patrick Suppes. 1960. *Axiomatic Set Theory*. Van Norstrand, Princeton.
- [Ta33] Alfred Tarski. 1933. *The concept of truth in the languages of the deductive sciences*. In *Logic, Semantics, Metamathematics, papers from 1923 to 1938* (p152-278). ed. John Corcoran. 1983. Hackett Publishing Company, Indianapolis.
- [Tu36] Alan Turing. 1936. *On computable numbers, with an application to the Entscheidungsproblem*. In M. Davis (ed.). 1965. *The Undecidable*. Raven Press, New York. Reprinted from the Proceedings of the London Mathematical Society, ser. 2. vol. 42 (1936-7), pp.230-265; corrections, Ibid, vol 43 (1937) pp. 544-546.
- [Wa63] Hao Wang. 1963. *A survey of Mathematical Logic*. North Holland Publishing Company, Amsterdam.
- [We27] Hermann Weyl. 1927. *Comments on Hilbert's second lecture on the foundations of mathematics*. In Jean van Heijenoort. 1967. Ed. *From Frege to Gödel: A source book in Mathematical Logic, 1878 - 1931*. Harvard University Press, Cambridge, Massachusetts.
- [An07a] Bhupinder Singh Anand. 2007. *The Mechanist's Challenge*. The Reasoner, Vol(1)5 p5-6.
- [An07b] 2007. *Why we shouldn't fault Lucas and Penrose for continuing to believe in the Gödelian argument against com*. The Reasoner, Vol(1)6 p3-4.
- [An07c] 2007. *Why we shouldn't fault Lucas and Penrose for continuing to believe in the Gödelian argument against com*. The Reasoner, Vol(1)7 p2-3.
- [An07d] 2007. *A constructive definition of the intuitive truth of the Axioms and Rules of Inference of Peano Arithmetic*. The Reasoner, Vol(1)8 p6-7.
- [An08] 2008. *Can we really falsify truth by dictat?*. The Reasoner, Vol(2)1 p7-8.

Authors postal address: 32 Agarwal House, D Road, Churchgate, Mumbai - 400 020, Maharashtra, India. Email: re@alixcomsi.com, anandb@vsnl.com.